BREATHLESSNESS

Dudgeon D, Shadd J. Assessment and management of dyspnea in palliative care. UpToDate 2015.

Jennings A-L, Davies AN, Higgins JPT, Gibbs JSR, Broadley KE. A systematic review of the use of opioids in the management of dyspnoea. Thorax.2002; 57: 939-944.

Abernethy AP, Currow DC, Frith P et al. Randomised, double blind, placebo controlled crossover trial of sustained release morphine for the management of refractory dyspnoea. BMJ.2003; 327: 523-26.

Bausewein C, Booth S, Gysels M, Higginson IJ. Non-pharmacological interventions for breathlessness in advanced stages of malignant non-malignant diseases. Cochrane Database of Systematic Reviews 2008, Issue 2. Art. No.: CD005623. DOI: 10.1002/14651858.pub2.

Galbraith S, Fagan P, Perkins P, Lynch A, Booth S. Does the use of a handheld fan improve chronic dyspnoea? A randomized, controlled, crossover trial. J Pain Symptom Manage. 2010; 39: 831-8.

Higginson I, Bausewein C, Reilly C et al. An integrated palliative and respiratory care service for patients with advanced disease and refractory breathlessness: a randomised controlled trial. The Lancet Respiratory Medicine. 2014; 2: 979-987.

Farquar M, Prevost AT, McCrone P et al. Is a specialist breathlessness service more effective and cost-effective for patients with advanced cancer and their carers than standard care? Findings of a mixed method randomised controlled trial. BMC Medicine. 2014; 12: 194.

Abernethy AP, McDonald CF, Frith PA et al. Effect of palliative oxygen versus room air in relief of breathlessness in patients with refractory dyspnoea: a double- blind, randomised controlled trial. Lancet. 2010; 376: 784-793.

Bradley JM, O'Neill BM. Short-term oxygen for chronic obstructive pulmonary disease. Cochrane Database of Systematic Reviews 2005.-issue 4. CD004356. DOI: 10.1002/14651858.CD004356.pub3.